

ORIGINAL ARTICLE



WILEY

Vitamin B5, a coenzyme A precursor, rescues TANGO2 deficiency disease-associated defects in *Drosophila* and human cells

Paria Asadi¹ | Miroslav P. Milev¹ | Djenann Saint-Dic¹ | Chiara Gamberi² | Michael Sacher^{1,3}

¹Department of Biology, Concordia University, Montreal, Quebec, Canada

²Department of Biology, Coastal Carolina University, Conway, South Carolina, USA

³Department of Anatomy and Cell Biology, McGill University, Montreal, Quebec, Canada

Correspondence

Michael Sacher, Department of Biology, Concordia University, Montreal, QC H4B1R6, Canada.
Email: michael.sacher@concordia.ca

Funding information

Canadian Institutes of Health Research; Natural Sciences and Engineering Research Council of Canada; TANGO2 Research Foundation; National Institute of Health (NIH) SCoRE, Grant/Award Number: P20GM103499-20; IDeA Network for Biomedical Research Excellence (INBRE) Developmental Research Program

Communicating Editor: Jerry Vockley

Abstract

Mutations in the Transport and Golgi Organization 2 (*TANGO2*) gene are associated with intellectual deficit, neurodevelopmental delay and regression. Individuals can also present with an acute metabolic crisis that includes rhabdomyolysis, cardiomyopathy, and cardiac arrhythmias, the latter of which are potentially lethal. While preventing metabolic crises has the potential to reduce mortality, no treatments currently exist for this condition. The function of *TANGO2* remains unknown but is suspected to be involved in some aspect of lipid metabolism. Here, we describe a model of *TANGO2*-related disease in the fruit fly *Drosophila melanogaster* that recapitulates crucial disease traits. Pairing a new fly model with human cells, we examined the effects of vitamin B5, a coenzyme A (CoA) precursor, on alleviating the cellular and organismal defects associated with *TANGO2* deficiency. We demonstrate that vitamin B5 specifically improves multiple defects associated with *TANGO2* loss-of-function in *Drosophila* and rescues membrane trafficking defects in human cells. We also observed a partial rescue of one of the fly defects by vitamin B3, though to a lesser extent than vitamin B5. Our data suggest that a B complex supplement containing vitamin B5/pantothenate may have therapeutic benefits in individuals with *TANGO2*-deficiency disease. Possible mechanisms for the rescue are discussed that may include restoration of lipid homeostasis.

KEYWORDS

coenzyme A, *Drosophila*, membrane traffic, metabolic crisis, neurodevelopment, *TANGO2*, vitamin B5

This study is dedicated to the memory of Dr. Nassim Shahrzad, an accomplished scientist with a bright future who was taken from her family, friends and colleagues much too soon. May the memory of her warm smile, collegial nature and devotion to her family serve as a source of comfort and inspiration to all those who knew her.

1 | INTRODUCTION

Initially identified nearly two decades ago as a protein that affects membrane trafficking in insect cells,¹ *TANGO2* is conserved from the fruit fly *Drosophila melanogaster* to humans and was only recently linked to human health.^{2,3}

Affected individuals with TANGO2-deficiency disease (TDD) manifest a plethora of clinical features (summarized in Schymick et al.⁴) including neurodevelopmental delay, seizures, developmental regression and intellectual deficit. Acute metabolic crises, often triggered by fever or fasting, manifest as hypoglycemia, lactic acidemia and elevated creatine phosphokinase and can lead to episodic rhabdomyolysis and potentially lethal cardiac arrhythmias.⁵ Little is known about the function and localization of the TANGO2 protein with studies suggesting mitochondrial, cytosolic or a mix of both localizations.^{2,3,6,7,8} Several reports have shown defective endomembrane trafficking and dilated endoplasmic reticulum (ER) in fibroblasts derived from TDD-affected individuals.^{3,7} Other studies have demonstrated defective mitochondrial morphology and respiration^{3,7,9} leading to the speculation that TANGO2 may affect some as yet unknown aspect of lipid metabolism. One study reported abnormal palmitate oxidation in cells derived from two individuals harboring two different TANGO2 alleles, suggesting a defect in fatty acid oxidation and/or oxidative phosphorylation.² These various results have made it difficult to devise treatment options for affected individuals, particularly to avoid the metabolic crisis which is the sole contributor of TDD lethality.

We hypothesized that if TANGO2 participates in lipid metabolism, then increasing the levels of the fatty acid activator Coenzyme A (CoA) might ameliorate cellular and organismal defects associated with TANGO2 loss-of-function. Pantothenic acid (also known as vitamin B5) is the obligate precursor of CoA in an essential pathway conserved in both prokaryotes and eukaryotes. In humans, the main source of pantothenic acid is dietary and a conserved, canonical five-step process converts this vitamin into CoA,¹⁰ though a possible parallel pathway has also been reported in *Drosophila*.¹¹ Both humans and *Drosophila* utilize four enzymes in the canonical pathway—three monofunctional enzymes (PANK, PPCS, and PPC-DC) that carry out the first three steps, and one bifunctional enzyme (COASY) catalyzing the final two steps¹⁰—further highlighting the conservation of this pathway. The generated CoA is used in a variety of cellular processes including the TCA cycle, lipid activation for fatty acid oxidation and synthesis as well as lipoylation of specific proteins. Some of these cellular processes are carried out by holo-acyl carrier proteins (holoACP), which themselves require the 4'-phosphopantetheine group from CoA for their production.¹² To test our hypothesis, we subjected two models of TDD to vitamin B5 supplementation and assessed phenotypic rescue. We report that the *Drosophila* TANGO2^{G517} loss-of-function mutants displayed several defects that overlap with the clinical phenotypes of TDD-affected individuals and can be used as a miniature whole-animal model of TANGO2 loss-of-function. We found that vitamin B5 supplementation rescued

the defective phenotypes of the TANGO2^{G517} mutant up to near wild type levels. These effects are specific to vitamin B5 since vitamins C, B3, and B9 supplementation did not result in a similar level of rescue, though vitamin B3 did show a modest effect in one assay. Continuous supplementation from larval development to adults was not required for rescue but positive effects appeared time-dependent. Similarly, in cultured human fibroblasts, vitamin B5 rescued a TANGO2-dependent cellular trafficking defect. Our results suggest that supplementation with a multivitamin or a B complex vitamin containing vitamin B5, and perhaps vitamin B3, may benefit individuals affected by TDD.

2 | METHODS

2.1 | *Drosophila melanogaster* maintenance, growth and vitamin supplementation

TANGO2^{G517} flies were obtained from the Bloomington *Drosophila* Stock Center. Control Oregon^R (Ore^R) and TANGO2^{G517} fruit flies were maintained at 20–23°C on a Bloomington formula commercial fly food (Genesee Scientific). Flies aged 0–2 days old were collected and aged 10 days to yield the 10–12 days old populations to assay. Flies were kept well-fed by transferring to fresh food every 3 days. Supplemental vitamin B5 (Sigma, P5155) (2–4 mM as indicated),¹¹ vitamin B3 (Sigma, N0636) (0.5 mM),¹³ vitamin B9 (Sigma, F7876) (10 µM),¹⁴ or vitamin C (ThermoFisher, A61-100) (7 mM)¹⁵ were dissolved in water and added to the warm liquid food and allowed to solidify at room temperature. Unused food was stored at room temperature for no longer than 10 days before use.

2.2 | Real time quantitative PCR (RT-qPCR)

Total RNA was collected from 50 to 200 fly heads using TRIzolTM reagent (ThermoFisher) as per the manufacturer's protocol. Extracted RNA was converted to cDNA using the First Strand cDNA Synthesis kit (Origene) and then subjected to real time quantitative PCR using SYBR Green Master mix (BioRad) and the following oligonucleotides: TANGO2-forward 5'-GGCTGCTCAAAGATTGCACC-3'; TANGO2-reverse 5'-TTGCCAAATCCGTAGCACTCG-3'; Rpl32-forward 5'-AGCATACAGGCCCAAGATCG-3'; Rpl32-reverse 5'-TGTTGTCGATACCCTTGGGC-3'. The experiment was performed three times, with each cDNA sample amplified in triplicate. The TANGO2 expression was normalized to that of Rpl32.

2.3 | *Drosophila* assays

All assays were performed on larvae and on both male and female adults (10–12 days old) of the *Ore^R* and *TANGO^{G517}* genotypes. The fly *TANGO2* gene is on the X chromosome, and we noticed that there were much fewer males in the *TANGO2^{G517}* colony. The reason for this remains unclear. Recovering higher numbers of female flies, assays were first performed with females and are shown in the main results section. Assays were then repeated with males and are shown in the supplemental information section. All assays were repeated a minimum of three times. Adults were anesthetized with CO₂ to separate males from females and allowed to reacclimate for 45 min prior to any assay. The total number of flies assayed from all biological replicates (*N*) is specified for each assay below. The *N* values for the specific experiments shown in each figure are indicated in the figure legends. The largest values represent controls which were repeated multiple times while testing various vitamin conditions. In each figure, one representative experiment is shown except for Figures 1D and 2B–D (and Figures S1D and S2B,C), for which cumulative results are displayed.

2.3.1 | Starvation assay

Flies of age 10–12 days old were placed into empty plastic vials and starved for food and fluids. Survival was monitored hourly in three assay replicates. The total *N* values ranged from 750 to 3000.

2.3.2 | Larvae motility assay

Third instar larvae were placed at the center of a 10 cm petri dish containing 2% agar in water. Larvae were observed over 9 min and the total crawling distance moved was recorded. The direction of movement was assessed by comparing the total distance moved from the initial position of larva on the dish (*d*) to the greatest displacement in any direction from the initial position (ΔX_{\max}). The total *N* values ranged from 15 to 60.

2.3.3 | Climbing (geotaxis) assay

The protocol of Tapia et al¹⁶ was modified as follows. The target line was placed 5 cm from the bottom of the tube and flies were given 10 s to reach this target line. The total *N* values ranged from 225 to 2500.

2.3.4 | Open field assay

The protocol of Tapia et al¹⁶ was followed except that the distance moved was recorded manually. Using a 10 cm diameter dish, the field was set as a 5 cm diameter from the center of the dish. The total *N* values ranged from 15 to 20.

2.3.5 | Learning assay

Flies were placed on 1% agarose in water for 16 h before acclimating for 10 min in the dark. The protocol of Tapia et al¹⁶ was modified as follows. A disk of filter paper wetted in 0.1 M quinine (a bitter substance that repulses flies) was placed in the lighted end of the setup, and the flies were allowed 30 s to venture into the lighted end of the setup. The total *N* values ranged from 40 to 50.

2.3.6 | Seizure induction

Flies were acclimated in an empty plastic vial for 45 min prior to the assay. The vials were then immersed in a beaker containing water at either 42°C (males) or 42.3°C (females) for 120 s and the time to the first seizure was noted. Flies counted to be experiencing a seizure were those who fell to the bottom of the tube and underwent random spasms. The total *N* values ranged from 225 to 300.

Statistical analyses were performed using GraphPad Prism as one-way ANOVA with a post hoc Tukey HSD test. Data are shown with *SEM* error bars.

2.4 | Membrane trafficking assay

The retention using selective hooks (RUSH) assay was performed as previously described.⁷ Human fibroblasts grown in DMEM with 10% fetal bovine serum were treated with 2 mM vitamin B5, 0.5 mM vitamin B3, or 0.5 mM vitamin C for 4 days prior to performing the assay. The assay was performed a minimum of three times for each condition.

3 | RESULTS

3.1 | *TANGO2^{G517}* mutant flies display multiple behavioral defects

The human (276 amino acids) and *Drosophila* (283 amino acids) *TANGO2* proteins have 30% identity and 50% similarity over their entire length. The predicted structures based on alphaFold¹⁷ are also similar, with the pairwise alignment giving a root mean square distance value of




FIGURE 1 The *Drosophila* *TANGO2*^{G517} mutants display behavioral defects that are rescued by vitamin B5 supplementation. (A) Female fly populations were placed into empty plastic vials and survival was examined every hour until all flies were dead. (B, C) Female fly populations were subjected to elevated temperatures for 120 s as detailed in the Methods section. The percentage of flies experiencing a seizure (B) and the time to the first seizure (C) are shown. (D) Female fly populations were subjected to a learning assay as described in the Methods section. The percentage of flies that learned to avoid the quinine-laced light side of the setup was recorded. Where indicated, supplements of either vitamin B5 or B3 were included in the food for at least one generation prior to the assays. The concentrations of vitamin B5 used were 4 mM in (A) and 2 mM in (B, C, and (D)). *N* values for each experiment shown were 75 in (A, B, and C), and 30–40 in (D). The asterisks for statistical analyses indicate **p* ≤ 0.05, ***p* ≤ 0.01, and *****p* ≤ 0.0001. CTRL represents *Ore^R* flies and TM represents *TANGO2*^{G517} flies. CTRL represents *Ore^R* flies and TM represents *TANGO2*^{G517} flies. Results obtained with male flies are shown in Figure S1.

1.83 and a template modeling score of 0.9 over 96% coverage. We therefore evaluated *Drosophila* as a model organism to study TDD. The *TANGO2*^{G517} allele has a P-element construct inserted at the 5' end of the *TANGO2* gene. Although these flies are viable, they are delayed in development relative to the *Ore^R* wild type control. *TANGO2* expression was measured by quantitative real-time PCR from RNA extracted from heads, which showed that the *TANGO2*^{G517} flies had a residual 3.13% ± 1.23 expression compared to that of wild type *Ore^R* control flies of equal age. Thus, the *TANGO2*^{G517} mutant flies can serve as a good model to examine the effects of *TANGO2* loss-of-function.

We initially performed a series of assays that replicate triggers of metabolic crises in TDD individuals and assays that measure some of the common non-neuromuscular

defects. Since starvation triggers a metabolic crisis in TDD individuals,¹⁸ we first examined the viability of 10–12 days old adult flies upon starvation. Both *Ore^R* and *TANGO2*^{G517} flies were starved for both food and moisture by placing them in empty vials. The *TANGO2*^{G517} flies died faster than *Ore^R* wild type flies with 50% viability reached at ~12 h for the former and ~25 h for the latter (Figures 1A and S1A).

A common feature of individuals suffering from TDD is seizures. We therefore assessed the heat-induced seizure rate of *TANGO2*^{G517} flies compared to *Ore^R* wild type. While the *Ore^R* wild types did not experience seizures due to the elevated temperatures, ~40% of *TANGO2*^{G517} flies did (Figures 1B and S1B), with the first seizure occurring at ~30 s (Figures 1C and S1C). Female flies required slightly higher temperatures (42.3°C) in




FIGURE 2 Motor defects associated with the *TANGO2*^{G517} mutants are rescued by vitamin B5 supplementation. (A) Female flies were placed into an empty vial, gently knocked to the bottom of the vial and were examined as they climbed. The percentages of flies that reached a line 5 cm from the bottom of the vial within 10 s are shown. (B) Third instar larvae were observed as they crawled over 9 min on a dish containing 2% agar. The total distance moved is shown. (C) Adult females were placed in the center of a 10 cm diameter petri dish and the distance moved over 10 min is displayed, as well as the time spent in the central area of the dish (D). Vitamins B5, B3, B9, or C as indicated were included in the food for at least one generation prior to the assays. The concentrations of vitamin B5 used were 4 mM in (A, C, and D), and 2 mM in (B). *N* values for each experiment shown were 50–75 in (A), 15–35 in (B) and 15 in (C, D). The asterisks for statistical analyses indicate $**p \leq 0.01$ and $****p \leq 0.0001$. CTRL represents *Ore*^R flies and TM represents *TANGO2*^{G517} flies. Results for male flies are shown in Figure S2.

order to induce seizures in the *TANGO2*^{G517} flies compared to males (42°C).

Since intellectual deficit is a common feature of TDD, we assessed the fly learning ability.^{16,19} In this assay, flies are kept in the dark and then allowed to move toward a bright area of the housing vial following their natural attraction to light. The bright end of the vial was, however, laced with quinine, a bitter substance repulsive to flies. The flies were then reacclimated for 10 s on the dark side of the vial and allowed to venture into the light side six more times before finally testing their ability to

avoid the quinine-laced side over a 10 s period. While ~65% of the *Ore*^R flies learned to avoid the repulsive lighted side, only about 30% of the female *TANGO2*^{G517} flies did (Figures 1D and S1D). We also noted that *Ore*^R flies already showed signs of learning during early phases of the training sessions (trials 1–5) prior to the actual test (trial 7) whereas it took longer for the *TANGO2*^{G517} flies to achieve avoidance (Figure S1E,F).

We next performed a series of assays to test possible motor defects. First, we examined geotaxis of the 10–12-days old adults by gently jolting them to the bottom of

an empty vial and assessing the rate at which they climb 5 cm to a target line. As above, the *TANGO2^{G517}* flies performed worse than control, with an ~80% reduction of climbing compared to *Ore^R* (Figures 2A and S2A).

We then examined third instar larvae for their ability to move on agar medium over 9 min. As was seen with the adults, the *TANGO2^{G517}* larvae displayed reduced movement by greater than 50% compared to *Ore^R* wild type (Figure 2B). In addition, while the wild type larvae tended to move in a straight line, the *TANGO2^{G517}* larvae moved in a more circular, and seemingly uncoordinated manner. This may be reminiscent of a form of ataxia often seen in individuals with TDD.⁴ In order to quantify this phenotype, we reasoned that larvae that move in a straight line will have a “total distance moved/maximum magnitude of displacement from the initial point” ($d/\Delta x_{\max}$) ratio of ~1. Mutant larvae with circular motions would remain closer to the initial point despite the total distance moved. Hence, $d/\Delta x_{\max}$ would be >1. Indeed, we found that *Ore^R* larvae had a $d/\Delta x_{\max}$ ratio of 1.036 ± 0.006 while *TANGO2^{G517}* larvae showed a significantly higher ratio of 1.378 ± 0.076 ($p < 0.0001$).

Open-field assays are used to assess anxiety in flies and other model organisms.^{16,20} We evaluated 10–12 days old flies for their ability to walk on a plastic culture dish and monitored their location either within a 5 cm diameter area from the center of the dish or peripherally to it. The *TANGO2^{G517}* flies walked on average ~75% shorter distances over the given time period compared to *Ore^R* wild type (Figures 2C and S2B). Interestingly, the *TANGO2^{G517}* flies showed a propensity to move toward the edge of the dish and linger, spending ~85% less time in the center of the dish than the *Ore^R* wild type (Figures 2D and S2C). Therefore, *TANGO2^{G517}* may exhibit anxious behavior.

3.2 | Vitamin B5 reverses all of the behavioral defects in *TANGO2^{G517}* flies

Having observed a number of defects reminiscent of TDD in *TANGO2^{G517}* mutants, and since *TANGO2* defects are thought to impair lipid metabolism, we sought to determine possible benefits of supporting lipid metabolism via vitamin B5 supplementation since this molecule is a precursor to the lipid activator CoA. We first titrated vitamin B5 levels, identifying 8 mM as the highest concentration that did not appear to negatively impact fly cultures (not shown). This amount is similar to that used in another study.¹¹ Using the relatively fast geotaxis assay (Figures 2A and S2A), we found that 2–4 mM vitamin B5 gave more robust results and settled on this range for all of the rescue experiments described below. In all cases, we found that vitamin B5 supplementation improved

assay performance of the *TANGO2^{G517}* flies. Compared to untreated siblings, the female 10–12 days old *TANGO2^{G517}* flies fed vitamin B5 were able to survive longer upon starvation (50% survival at ~25 h compared to 12 h for vehicle-treated sibling flies) (Figure 1A), had a significant reduction (~95% reduction) of heat-induced seizures and ~3.5-fold longer times before experiencing the first seizure (Figures 1B,C), performed nearly 2-fold better in the learning assay (Figures 1D and S1F), had ~3.5-fold stronger geotactic response (Figure 2A), walked ~3-fold further (Figure 2C), and stayed in the center of the culture dish ~3-fold longer (Figure 2D). Similar effects of vitamin B5 were seen in the male flies (Figures S1A–E, S2A–C). The third instar larvae also showed increased movement after vitamin B5 supplementation compared to untreated larvae (Figure 2B), and regained the ability to move in a straight line rather than the uncoordinated circular movement typical of untreated siblings ($d/\Delta x_{\max}$ ratio of 1.120 ± 0.022 for vitamin B5-treated larvae compared to 1.378 ± 0.076 for untreated larvae [$p < 0.001$]).

To determine if these effects were specific to vitamin B5, we tested other vitamins including two other B vitamins (B3 and B9) as well as vitamin C. While vitamins B9 and C did not improve larval crawling, seizures, and open field tests respectively (Figures 1B,C and 2B–D), vitamin B3 displayed a small but significant improvement of motor activity in the geotaxis assay (Figure 2A).

3.3 | Vitamin B5 shows a time-dependent rescue of behavioral defects

In the previous assays, flies were receiving vitamin B5 supplementation throughout development from the egg stage until 10–12 days old when adults were assayed. We next asked if adult *TANGO2^{G517}* flies grown in normal conditions could benefit from vitamin B5 supplementation over shorter periods of time. To address this, we used the geotaxis assay, the starvation assay, and the heat-induced seizure assay that recapitulate critical features of TDD. The *TANGO2^{G517}* flies tested were fed either regular food (untreated) or food supplemented with vitamin B5 since the egg stage until the 10–12 days old age when adults were assayed, or grown in regular food and fed vitamin B5-supplemented food 1, 2, or 3 days prior to the assay. The *TANGO2^{G517}* flies showed a time-dependent increase in geotaxis (Figures 3A and S3A) and survival upon starvation (Figure 3B), and reduced incidence of heat-induced seizures upon vitamin B5 supplementation (Figures 3C and S3B). In each case, growth on vitamin B5 for as little as 1 day significantly rescued the *TANGO2^{G517}* defects, suggesting that although prolonged




FIGURE 3 Vitamin B5 supplementation shows a time-dependent rescue of phenotypes in the *TANGO2*^{G517} mutants. *TANGO2*^{G517} female flies were grown either on food without vitamin B5, or grown for at least one generation either on regular food or on vitamin B5-supplemented food. A portion of the untreated *TANGO2*^{G517} adult flies were placed on vitamin B5-containing food either 1, 2, or 3 days prior to the assays. Flies were then assayed for their climbing ability (A), survival during starvation (B) and resistance to heat-induced seizures (C). The concentrations of vitamin B5 used were 4 mM in (A) and (C), and 2 mM in (B). *N* values for each experiment shown were 50–75 in (A), 75 in (B) and 75 in (C). The asterisks for statistical analyses indicate **p* ≤ 0.05, ***p* ≤ 0.01, ****p* ≤ 0.001, and *****p* ≤ 0.0001. TM represents *TANGO2*^{G517} flies. Results obtained with male flies are shown in Figure S3. Note that although blue coloring has been used for vitamin B5 supplementation in all figures, different coloring has been used in panel B to ensure ease of visualization of the curves.

administration may prove most effective, vitamin B5 may exert benefits over a relatively short time frame.

3.4 | Vitamin B5 rescues a membrane trafficking defect in human fibroblasts devoid of TANGO2

Given the wide range of rescuing effects of vitamin B5 in the TDD *Drosophila* model, we asked if similar benefits would be found in mammals. We examined fibroblasts from a compound heterozygous individual with the common *TANGO2* exon 3–9 deletion and *TANGO2* exon

6 deletion, effectively rendering the cells a *TANGO2* knockout,^{2,7} and compared them to control fibroblasts. We first empirically determined that 2 mM vitamin B5 and 0.5 mM vitamin B3 supplementation in the growth medium for up to 4 days had no negative effects on the growth or morphology of the fibroblasts (not shown). Using an established assay for ER to Golgi transport, and as previously reported,⁷ we found the fibroblasts with the *TANGO2* alleles to have a greatly reduced rate of transport of the cargo protein from the ER to the Golgi (Figure 4A). Following vitamin B5 supplementation, the rate of transport was greatly increased to near control levels. The level of rescue was similar to that seen when




FIGURE 4 Vitamin B5 rescues an endoplasmic reticulum-to-Golgi trafficking defect in *TANGO2*-deficient human fibroblasts. (A) Wild type (CTRL) fibroblasts or fibroblasts harboring the compound heterozygous exon 3-9 del/exon 6 del allele (*TANGO2*-deficient cells) were transfected with a plasmid expressing sialyltransferase-GFP-streptavidin binding protein which is retained in the ER by association with an ER-localized hook composed of streptavidin. After release of the GFP-tagged protein by biotin, fluorescence was monitored in the Golgi over time and quantified. Solid lines represent CTRL cells and dashed lines represent the *TANGO2* mutant cells. Black coloring represents untreated cells, red coloring represents vitamin B5-treated cells, blue coloring represents vitamin B3-treated cells, gray coloring represents vitamin B9-treated cells and green coloring represents vitamin C-treated cells. Note that the only dashed line that clusters with the solid lines is from the vitamin B5-treated cells. (B) Lysates from CTRL and *TANGO2*-deficient cells either untreated or treated with vitamin B5 for 4 days were probed for *TANGO2* and tubulin as a loading control. Note that the protein product of the exon 6 del allele would be expected to appear at ~15 kD if it was produced and stable.

the cells were rescued with wild type *TANGO2*,⁷ suggesting that vitamin B5 could functionally replace *TANGO2* in this assay. In these cells one allele (exon 3-9 del) is effectively a *TANGO2* knockout, while the other allele (exon 6 del) could theoretically produce a truncated protein of ~15 kD. Thus, it was formally possible that vitamin B5 could increase expression of the ~15 kD truncation which might explain the rescue observed. However, as shown in Figure 4B, vitamin B5 treatment of the *TANGO2* deficient fibroblasts did not result in the appearance of truncated *TANGO2* protein. This suggests that the mechanism of action of vitamin B5, at least with respect to *TANGO2*, does not seem to involve an induction in protein expression. Corroborating the conclusion that vitamin B5 specifically rescues the trafficking defect

of *TANGO2* deficiency, neither vitamin B3, B9 nor C supplementation showed a similar augmented rate of trafficking (Figure 4A). Collectively, our study shows that vitamin B5 supplementation can rescue a wide range of defects associated with *TANGO2* loss-of-function in *TANGO2*-deficient cells and whole organisms from *Drosophila* to humans.

4 | DISCUSSION

TANGO2-deficiency disease is a multisystem organ disease that includes neuromuscular, developmental, and intellectual delays. Affected individuals are susceptible to clinical deterioration when exposed to metabolic stressors

such as fever or fasting. These episodes of metabolic crisis lead to worsening ataxia, muscle weakness, and cardiac arrhythmias as well as cardiomyopathy and are the leading cause of mortality in this disease. Here, we find that *Drosophila* *TANGO2*^{G517} mutants exhibit greatly reduced *TANGO2* expression and display defects that recapitulate many of these clinical features. Thus, these mutants may be used to model TDD. Interestingly, all of the phenotypic defects seen in the *TANGO2*^{G517} flies are significantly reversed, some to near wild type levels, by vitamin B5 supplementation. It is noteworthy that a motor function was also rescued, albeit to a much lower, but still significant, degree by vitamin B3 that was shown to have broad positive effects in several unrelated conditions.²¹

To date, there are no treatment options for individuals suffering from TDD. A recent study suggests that folic acid (vitamin B9) can restore normal rhythm in a patient-derived cardiomyocyte model system.²² Together with our present study this suggests that several B complex vitamins may have beneficial effects on various clinical aspects of this disease. Consistent with this notion, a natural history study of over 70 individuals suffering from TDD found that metabolic crises were not seen in individuals taking a multivitamin or a B complex supplement.²³ Multivitamins typically contain B vitamins including B5. Thus, our work and the aforementioned studies strongly suggest that B complex supplementation with at least vitamins B5 and B9, and possibly B3, may offer substantial therapeutic benefits for those affected by TDD.

The mechanism whereby these B vitamins rescue the *TANGO2*-dependent defects is currently unknown and awaits a thorough understanding of the molecular function(s) of the *TANGO2* protein. Based on the trafficking assay in human fibroblasts (Figure 4A) the most likely scenario is that production of a metabolite, and not increased production of the *TANGO2* protein, rescues the critical ER-to-Golgi function. In light of vitamin B5 being a precursor to CoA, it is tempting to speculate that increased levels of an unknown CoA metabolite(s) lead to a downstream phenotypic rescue in the two model systems.

The five-step biosynthesis pathway converting vitamin B5 to CoA has been linked to human neurodegenerative diseases with brain iron accumulation (NBIA). Variants in *PANK2*, mediating the first step in the pathway, and *COASY*, regulating the last two steps, are linked to pantothenate kinase associated neurodegeneration (PKAN or NBIA1; OMIM 234200) and *COASY* protein-associated neurodegeneration (CoPAN or NBIA6; OMIM 618266). Both conditions are characterized by impaired de novo synthesis of CoA. While individuals with PKAN, CoPAN and many with TDD

exhibit cognitive impairment and movement disorders, TDD-affected individuals do not demonstrate the pathognomonic finding of brain iron accumulation observed in PKAN and CoPAN individuals.

Based on these differences, we can speculate on several possible cellular processes downstream of CoA that may be affected in TDD individuals. Acyl-CoA acts as a source of fatty acid for β -oxidation in mitochondrial energy production as well as a substrate for protein acylation. The *TANGO2* protein was originally thought to function in membrane trafficking and it is noteworthy that this process is regulated by protein acylation²⁴ suggesting this as a possible affected process in TDD. Another possibility is that *TANGO2* acts downstream of holoACP, formed from apoACP by the transfer of the 4'-phosphopantetheine from CoA. holoACP is involved in fatty acid synthesis, defects in which could affect lipid homeostasis and the function of multiple membranes, a process that would respond to the dietary changes that trigger metabolic crises. Consistent with this notion are studies reporting minimal,¹⁸ modest,^{2,8} and strong²⁵ changes in acylcarnitines in TDD-affected individuals. Further supporting this notion are several recent studies that demonstrated profound changes in the cellular lipid profile in a *TANGO2* knockdown model which were further exaggerated upon starvation,²⁶ and in a zebrafish model,²⁷ consistent with preliminary lipidomic data from our group on fibroblasts devoid of *TANGO2*. holoACP is also involved in lipoylation of four protein complexes involved in mitochondrial and amino acid metabolism,²⁸ possibly explaining the reported oxidative phosphorylation defects and triggered metabolic crises. Alternatively, holoACP is involved in the synthesis of iron-sulfur clusters.²⁹ In this respect it is noteworthy that a recent study suggests *TANGO2* may act as a heme chaperone, moving this iron-containing molecule out of the mitochondria.³⁰ If so, it remains to be determined how a TDD trigger relates to heme transport and a metabolic crisis, and why brain iron accumulation is not seen in TDD affected individuals.

Vitamin B3 is used in a salvage pathway to generate nicotinamide adenine dinucleotide (NAD). The latter molecule is a cofactor required in a number of cellular processes, perhaps best known as an electron acceptor/donor in energy production in the mitochondria. Given that several studies have suggested that a portion of the *TANGO2* protein associates with mitochondria in humans and mice,^{7,9,31} and a mitochondrial defect may be a secondary consequence of *TANGO2* deficiency, the modest benefit of vitamin B3 in the climbing assay may reflect a nonspecific overall increase in energy production or other factors, such as mitigation of oxidative stress.

Consistent with this notion, vitamin B3 has been shown to be beneficial in a number of different, unrelated diseases.²¹

A precise definition of all TANGO2 cellular roles in the functional context of different tissues and organs will be needed to understand TDD and design targeted therapeutics. However, studies in several models support the conclusion that vitamins B5, B9, and perhaps B3 can alleviate and/or remove dangerous TDD manifestations even with a short course of administration. Considering the extensive knowledge about vitamin B supplementation and the minimal toxicity, these studies may improve the current care of individuals affected by TDD.

AUTHOR CONTRIBUTIONS

Paria Asadi performed and analyzed all of the *Drosophila* assays. Miroslav P. Milev performed and analyzed the human fibroblast data. Djenann Saint-Dic performed qPCR and analyzed the data. Chiara Gamberi conceived and guided the *Drosophila* work and analyzed the data. Michael Sacher conceived the study, guided the work, analyzed the data and wrote the manuscript. All authors edited the manuscript and approved the final version.

ACKNOWLEDGMENTS

We are grateful to members of the Sacher and Gamberi laboratories for critical review of this work, as well as to Drs. S. Mackenzie, L. Zhang, C. Miyake, S. Lalani, and L. Ghaloul-Gonzalez for many helpful discussions. Stocks obtained from the Bloomington *Drosophila* Stock Center (NIH P40D018537) were used in this study. Imaging was performed at the Centre for Microscopy and Cellular Imaging at Concordia University.

FUNDING INFORMATION

This work was supported by the Canadian Institutes of Health Research (Michael Sacher), the Natural Sciences and Engineering Research Council of Canada (Michael Sacher), the TANGO2 Research Foundation (Michael Sacher and Chiara Gamberi), the National Institute of Health (NIH) SCoRE grant P20GM103499-20 (Chiara Gamberi) and the IDeA Network for Biomedical Research Excellence (INBRE) Developmental Research Program grant (Chiara Gamberi). Chiara Gamberi is a member of the IDeA Networks of Biomedical Research Excellence (INBRE), the Center of Excellence in Research on Orphan Diseases—Fondation Courtois, and the Disease Modeling Research Center at Coastal Carolina University.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data supporting this study are available from the corresponding author upon reasonable request.

ETHICS STATEMENT

Work on human fibroblasts has been approved by the Research Ethics committee of Concordia University.

ORCID

Michael Sacher  <https://orcid.org/0000-0003-2926-5064>

REFERENCES







- Bard F, Casano L, Mallabiabarrena A, et al. Functional genomics reveals genes involved in protein secretion and Golgi organization. *Nature*. 2006;439:604-607.
- Kremer LS, Distelmaier F, Alhaddad B, et al. Bi-allelic truncating mutations in TANGO2 cause infancy-onset recurrent metabolic crises with encephalocardiomyopathy. *Am J Hum Genet*. 2016;98:358-362.
- Lalani SR, Liu P, Rosenfeld JA, et al. Recurrent muscle weakness with rhabdomyolysis, metabolic crises, and cardiac arrhythmia due to bi-allelic TANGO2 mutations. *Am J Hum Genet*. 2016;98:347-357.
- Schymick J, Leahy P, Cowan T, et al. Variable clinical severity in TANGO2 deficiency: case series and literature review. *Am J Med Genet A*. 2022;188:473-487.
- Miyake CY, Lay EJ, Beach CM, et al. Cardiac crises: cardiac arrhythmias and cardiomyopathy during TANGO2 deficiency related metabolic crises. *Heart Rhythm*. 2022;19:1673-1681.
- Jennions E, Hedberg-Oldfors C, Berglund AK, et al. TANGO2 deficiency as a cause of neurodevelopmental delay with indirect effects on mitochondrial energy metabolism. *J Inherit Metab Dis*. 2019;42:898-908.
- Milev MP, Saint-Dic D, Zardoui K, et al. The phenotype associated with variants in TANGO2 may be explained by a dual role of the protein in ER-to-Golgi transport and at the mitochondria. *J Inherit Metab Dis*. 2021;44:426-437.
- Mingirulli N, Pyle A, Hathazi D, et al. Clinical presentation and proteomic signature of patients with TANGO2 mutations. *J Inherit Metab Dis*. 2020;43:297-308.
- Heiman P, Mohsen AW, Karunanidhi A, et al. Mitochondrial dysfunction associated with TANGO2 deficiency. *Sci Rep*. 2022;12:3045.
- Leonardi R, Jackowski S. Biosynthesis of pantothenic acid and coenzyme A. *EcoSal Plus*. 2007;2:10.
- Rana A, Seinen E, Siudeja K, et al. Pantethine rescues a *drosophila* model for pantothenate kinase-associated neurodegeneration. *Proc Natl Acad Sci U S A*. 2010;107:6988-6993.
- Beld J, Sonnenschein EC, Vickery CR, Noel JP, Burkart MD. The phosphopantetheinyl transferases: catalysis of a post-translational modification crucial for life. *Nat Prod Rep*. 2014;31:61-108.
- Jia H, Li X, Gao H, et al. High doses of nicotinamide prevent oxidative mitochondrial dysfunction in a cellular model and improve motor deficit in a *drosophila* model of Parkinson's disease. *J Neurosci Res*. 2008;86:2083-2090.
- Blatch SA, Meyer KW, Harrison JF effects of dietary folic acid level and symbiotic folate production on fitness and




- development in the fruit fly *Drosophila melanogaster*. *Fly (Austin)*. 2010;4:312-319.
15. Bahadorani S, Bahadorani P, Phillips JP, Hilliker AJ. The effects of vitamin supplementation on drosophila life span under normoxia and under oxidative stress. *J Gerontol A Biol Sci Med Sci*. 2008;63:35-42.
 16. Tapia A, Giachello CN, Palomino-Schatzlein M, Baines RA, Galindo MI. Generation and characterization of the *Drosophila melanogaster* paralytic Gene Knock-out as a model for Dravet syndrome. *Life (Basel)*. 2021;11(11):1261.
 17. Jumper J, Evans R, Pritzel A, et al. Highly accurate protein structure prediction with AlphaFold. *Nature*. 2021;596:583-589.
 18. Berat CM, Montealegre S, Wiedemann A, et al. Clinical and biological characterization of 20 patients with TANGO2 deficiency indicates novel triggers of metabolic crises and no primary energetic defect. *J Inherit Metab Dis*. 2021;44:415-425.
 19. Ali YO, Escala W, Ruan K, Zhai RG. Assaying locomotor, learning, and memory deficits in drosophila models of neurodegeneration. *J Vis Exp*. 2011;49:2504.
 20. Napoli E, Song G, Liu S, et al. Zdhc13-dependent Drp1 S-palmitoylation impacts brain bioenergetics, anxiety, coordination and motor skills. *Sci Rep*. 2017;7:12796.
 21. Fricker RA, Green EL, Jenkins SI, Griffin SM. The influence of nicotinamide on health and disease in the central nervous system. *Int J Tryptophan Res*. 2018;11:1178646918776658.
 22. Zhang L, Xu W, Cao Y, et al. Folate ameliorates arrhythmia in patient-derived TANGO2-deficient iPSC-cardiomyocytes. *Res Square*. 2022. [10.21203/rs.3.rs-1778084/v1](https://doi.org/10.21203/rs.3.rs-1778084/v1)
 23. Miyake C, Lay E, Soler-Alfonso C, et al. Natural history of TANGO2 deficiency disorder: baseline assessment of 73 patients. *Genet Med*. 2022; (in press).
 24. Ernst AM, Toomre D, Bogan JS. Acylation - a new means to control traffic through the Golgi. *Front Cell Dev Biol*. 2019; 7:109.
 25. Yokoi K, Nakajima Y, Takahashi Y, et al. Transport and Golgi organization 2 deficiency with a prominent elevation of C14:1 during a metabolic crisis: a case report. *J Inherit Metab Dis Rep*. 2022; (in press).
 26. Lujan A, Foresti O, Brouwers N, et al. Defects in lipid homeostasis reflect the function of TANGO2 in acyl-CoA metabolism. *BioRxIV*. 2022. [10.1101/2022.11.05.515282](https://doi.org/10.1101/2022.11.05.515282)
 27. Casey J, Kim E, Tao B, et al. Glycerolipid defects in skeletal muscle contribute to rhabdomyolysis in Tango2 deficiency. *BioRxIV*. 2022. [10.1101/2022.11.12.516274](https://doi.org/10.1101/2022.11.12.516274)
 28. Rowland EA, Snowden CK, Cristea IM protein lipoylation: an evolutionarily conserved metabolic regulator of health and disease. *Curr Opin Chem Biol*. 2018;42:76-85.
 29. van Vranken JG, Jeong MY, Wei P, et al. The mitochondrial acyl carrier protein (ACP) coordinates mitochondrial fatty acid synthesis with iron sulfur cluster biogenesis. *Elife*. 2016;5:e17828.
 30. Sun F, Zhao Z, Willoughby MM, et al. HRG-9 homologues regulate haem trafficking from haem-enriched compartments. *Nature*. 2022;610:768-774.
 31. Maynard TM, Meechan DW, Dudevoir ML, et al. Mitochondrial localization and function of a subset of 22q11 deletion syndrome candidate genes. *Mol Cell Neurosci*. 2008;39:439-451.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.


How to cite this article: Asadi P, Milev MP, Saint-Dic D, Gamberi C, Sacher M. Vitamin B5, a coenzyme A precursor, rescues TANGO2 deficiency disease-associated defects in *Drosophila* and human cells. *J Inherit Metab Dis*. 2022;1-11. doi:[10.1002/jimd.12579](https://doi.org/10.1002/jimd.12579)

A**B****C****D****E****F**

A**B****C**

A

% flies above target line in 10 sec

**B**

% of flies with seizure

